

2008

The haemolytic effect of verapamil on erythrocytes exposed to varying osmolarity


Watts, T.

Watts, T. and Handy, R. (2008) 'The haemolytic effect of verapamil on erythrocytes exposed to varying osmolarity', The Plymouth Student Scientist, 1(2), p. 355.

<http://hdl.handle.net/10026.1/13822>

The Plymouth Student Scientist
University of Plymouth

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.



The haemolytic effect of verapamil on erythrocytes exposed to varying osmolarity

Tim Watts & Richard Handy

2008

Project Advisor: [Richard Handy](#), School of Biological Sciences, University of Plymouth, Drake Circus, Plymouth, PL4 8AA

Abstract

The haemolytic effect of verapamil on red blood cells (RBCs) exposed to varying osmolarity was investigated. The experimental approach used a modified red cell haemolysis assay with concentrations of verapamil ranging from 50–1500 IM compared to drug free controls. The time-course of haemolytic effects was also investigated. We also briefly determined the haemolytic effects of verapamil in Ca²⁺-free conditions (with added EGTA). In conditions representing decreasing osmolarity (dilution from 140–0 mM NaCl) there was a significant increase in erythrocyte haemolysis that was also dependent on verapamil concentration (ANOVA, $p < 0.05$). The red cells also showed a significantly increased rate of haemolysis over 5 h with increasing verapamil concentration (ANOVA, $p < 0.05$). The degree of RBC hypotonic haemolysis was significantly increased in a Ca²⁺-free medium (+EGTA) compared to normal saline and this effect was exacerbated by additions of verapamil (ANOVA, $p < 0.05$). Overall the data suggested that verapamil can cause haemolysis of RBCs in a predictable time- and concentration-dependent manner, and that verapamil increases the fragility of the erythrocytes further during hypotonic osmotic stress and Ca²⁺-free conditions. The mechanism of verapamil-dependent haemolysis could be directly related to the observed biphasic concentration-effect and could consequently involve several ion transport pathways.

Keywords: Verapamil; Haemolysis; Equine red blood cell; Calcium; Osmotic stress

[Click here to access the full article via ScienceDirect](#)